

Supplemental Materials
for
“Time-series Analysis of Mortality Effects of
Fine Particulate Matter Components in Detroit and Seattle”

Jiang Zhou, Kazuhiko Ito, Ramona Lall, Morton Lippmann, and George Thurston
New York University School of Medicine, Tuxedo, NY

Table of contents

Supplemental Material Figure 1: Excess mortality risk in the warm season in Detroit.


Supplemental Material Figure 2: Excess mortality risk in the cold season in Detroit.

Supplemental Material Figure 3: Excess mortality risk in the warm season in Seattle.


Supplemental Material Figure 4: Excess mortality risk in the cold season in Seattle.

Supplemental Material Figure 5: All cause mortality effect sensitivity analysis in Detroit.


Supplemental Material Figure 6: All cause mortality effect sensitivity analysis in Seattle.




Supplemental Material Figure 1: Excess mortality risk in the warm season in Detroit. The diamonds and triangles represent significantly ($p < 0.05$) positive and negative associations, respectively









Supplemental Material Figure 2: Excess mortality risk in the cold season in Detroit. The diamonds and triangles represent significantly ($p < 0.05$) positive and negative associations, respectively









Supplemental Material Figure 3: Excess mortality risk in the warm season in Seattle. The diamonds and triangles represent significantly ($p < 0.05$) positive and negative associations, respectively



Supplemental Material Figure 4: Excess mortality risk in the cold season in Seattle. The diamonds and triangles represent significantly ($p < 0.05$) positive and negative associations, respectively



Supplemental Material Figure 5: All cause mortality effect sensitivity analysis in Detroit, alternative degrees of freedom per year (5 and 12 df/year) compared to original degrees of freedom per year (8 df/year). Each point represents a pair of risk estimates for an air pollutant at a given lag evaluated by the two models.



Supplemental Material Figure 6: All cause mortality effect sensitivity analysis in Seattle, alternative degrees of freedom per year (5 and 12 df/year) compared to original degrees of freedom per year (8 df/year). Each point represents a pair of risk estimates for an air pollutant at a given lag evaluated by the two models.